

06-08-00  
FISH & RICHARDSON P.C.

A  
45 Rockefeller Plaza  
Suite 2800  
New York, New York  
10111

Telephone  
212 765-5070

Facsimile  
212 258-2291

Web Site  
www.fr.com

jc571 U.S. PTO  
09/589419  
06/07/00

June 7, 2000

Attorney Docket No.: 11127-002002

**Box Patent Application**  
Commissioner for Patents  
Washington, DC 20231

Presented for filing is a new divisional patent application of:

Applicant: TADASHI KOHNO, ISABURO HOSOI, JUNKO OHSHIMA,  
KUNIIHIKO SHIBATA AND ASUKA ITO

Title: DIAGNOSTIC AGENTS FOR PANCREATIC EXOCRINE  
FUNCTION

The prior application is assigned of record to Tokyo Gas Company Limited,  
a Japan corporation, by virtue of an assignment submitted to the Patent and  
Trademark Office for recording on September 23, 1999 at 010271/0420.

Enclosed are the following papers, including those required to receive a filing date  
under 37 CFR 1.53(b):

	<u>Pages</u>
Specification	39
Claims	6
Abstract	1
Declaration	6
Drawing(s)	11

Enclosures:

— Form PTO-1449, 2 pages, listing documents cited in the parent  
application(s). Please confirm that these have been considered in this

CERTIFICATE OF MAILING BY EXPRESS MAIL

Express Mail Label No. EE6472811194US

I hereby certify under 37 CFR §1.10 that this correspondence is being  
deposited with the United States Postal Service as Express Mail Post  
Office to Addressee with sufficient postage on the date indicated below  
and is addressed to the Commissioner for Patents, Washington,  
D.C. 20231

Date of Deposit June 7, 2000

Signature *Valentin Figueroa*

VALENTIN FIGUEROA  
Typed or Printed Name of Person Signing Certificate

06/07/00  
jc813 U.S. PTO  
Frederick P. Fish  
1855-1930  
W.K. Richardson  
1859-1951



BOSTON

DALLAS

DELAWARE

NEW YORK

SAN DIEGO

SILICON VALLEY

TWIN CITIES

WASHINGTON, DC

06/07/00  
jc571 U.S. PTO  
09/589419

FISH & RICHARDSON P.C.

Commissioner for Patents

June 7, 2000

Page 2

application by returning a copy of the Form PTO-1449 with the examiner's initials.

- Preliminary amendment, 5 pages.
- Postcard.

This application is a divisional (and claims the benefit of priority under 35 USC 120) of U.S. application serial no. 09/401,739, filed September 23, 1999. The disclosure of the prior application is considered part of (and is incorporated by reference in) the disclosure of this application.

Basic filing fee	\$690
Total claims in excess of 20 times \$18	\$72
Independent claims in excess of 3 times \$78	\$78
Fee for multiple dependent claims	\$0
Total filing fee:	\$840

A check for the filing fee is enclosed. Please apply any other required fees or any credits to deposit account 06-1050, referencing the attorney docket number shown above.


If this application is found to be incomplete, or if a telephone conference would otherwise be helpful, please call the undersigned at (212) 765-5070.

Kindly acknowledge receipt of this application by returning the enclosed postcard.

Please send all correspondence to:

CHRIS T. MIZUMOTO  
Fish & Richardson P.C.  
45 Rockefeller Plaza, Suite 2800  
New York, NY 10111

Respectfully submitted,



Chris T. Mizumoto  
Reg. No. 42,899  
Enclosures  
CTM/rap  
30019256 doc

007050" 6745450

06/07/00  
jc813 U.S. PTO

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant : Kohno et al.  
Serial No. :  
Filed : June 7, 2000  
Title : DIAGNOSTIC AGENTS FOR PANCREATIC EXOCRINE FUNCTION

Art Unit : Unknown  
Examiner : Unknown

**BOX PATENT APPLICATION**

Commissioner for Patents  
Washington, D.C. 20231

PRELIMINARY AMENDMENT

Prior to examination, please amend the application as follows:

In the Specification:

On page 1, line 5, after "Background of the Invention" insert – This application is a division of U.S. application serial no. 09/401,739, filed September 23, 1999, which--.

In the Claims:

Please cancel claims 25-42.

Claims 1-24 are left pending. The non-amended claims are reproduced in small print for ease of reference. For the calculation of the claims fees, please refer to the claims of this preliminary amendment. All of the multiply dependent claims have been amended to become singly dependent claims.

1. A  $^{13}\text{C}$ -labeled oligosaccharide or polysaccharide or a salt thereof excluding  $\text{U-}^{13}\text{C}$ -maltose,  $^{13}\text{C}$ -starch,  $1\text{-}^{13}\text{C}$ -maltotetraose and  $1\text{-}^{13}\text{C}$ -amylose.

2. The  $^{13}\text{C}$ -labeled oligosaccharide or polysaccharide or salt thereof according to claim 1, which is hydrolyzed with  $\alpha$ -amylase.

CERTIFICATE OF MAILING BY FIRST CLASS MAIL

I hereby certify under 37 CFR §1.8(a) that this correspondence is being deposited with the United States Postal Service as first class mail with sufficient postage on the date indicated below and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231.

Date of Deposit June 7, 2000  
Signature Valentin Figueroa  
Typed or Printed Name of Person Signing Certificate VALENTIN FIGUEROA

09589419-060700

12. The inclusion complex or salt thereof according to claim 11, wherein the host molecule is <sup>13</sup>C-labeled.

13. The inclusion complex or salt thereof according to claim 11, wherein the guest molecule is  $^{13}\text{C}$ -labeled.

14. (Amended) The inclusion complex or salt thereof according to [any one of claims 11 to 13] claim 11, wherein the guest molecule is selected from the group consisting of oligosaccharides, amino acids, peptides, organic acids, fatty acids, fatty acid glycerides, vitamins, catechins, carotinoids, flavonoids, and cholesterol.

15. The inclusion complex or salt thereof according to claim 14, wherein the guest molecule is selected from the group consisting of  $^{13}\text{C}$ -phenylalanine, benzoylphenylalanyl- $^{13}\text{C}$ -leucine and benzoylphenylalanyl- $^{13}\text{C}$ -leucine methyl ester.

16. (Amended) [A diagnostic agent for] A method for diagnosing pancreatic exocrine function, comprising administering a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled oligosaccharide or polysaccharide or a salt thereof or a derivative thereof other than  $^{13}\text{C}$ -starch to a subject.

17. (Amended) [The diagnostic agent for pancreatic exocrine function according to claim 16] The method according to claim 16, wherein the  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled oligosaccharide or polysaccharide or salt thereof or derivative thereof is hydrolyzed with  $\alpha$ -amylase.

18. (Amended) [The diagnostic agent for pancreatic exocrine function according to claim 17] The method according to claim 17, wherein the  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled oligosaccharide or polysaccharide or salt thereof or derivative thereof is not hydrolyzed with  $\alpha$ -glucosidase.

19. (Amended) [The diagnostic agent for pancreatic exocrine function according to any one of claims 16 to 18] The method according to claim 16, wherein the  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled oligosaccharide or polysaccharide is a linear or branched oligosaccharide or polysaccharide.

20. (Amended) [The diagnostic agent for pancreatic exocrine function according to claim 19] The method according to claim 19, wherein the  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled oligosaccharide or polysaccharide is modified at the non-reducing terminal.

095894.9-060700

21. (Amended) [The diagnostic agent for pancreatic exocrine function according to any one of claims 16 to 18] The method according to claim 16, wherein the  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled oligosaccharide or polysaccharide is a cyclic oligosaccharide or polysaccharide.

22. (Amended) [A diagnostic agent for] A method for diagnosing pancreatic exocrine function, comprising administering to a subject a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled inclusion complex or a salt thereof having a cyclodextrin or a modified derivative thereof as a host molecule.

23. (Amended) [The diagnostic agent for pancreatic exocrine function according to any one of claims 16 to 22] The method according to claim 16, wherein the pancreatic exocrine function to be diagnosed is the ability of the pancreas to secrete  $\alpha$ -amylase.

24. (Amended) [The diagnostic agent for pancreatic exocrine function according to any one of claims 16 to 22] The method according to claim 16, wherein the pancreatic exocrine function to be diagnosed is the ability of the pancreas to secrete  $\alpha$ -amylase and at least one pancreatic exocrine enzyme other than  $\alpha$ -amylase.

#### REMARKS

By this amendment, claims 25-42 have been cancelled. Accordingly, claims 1-24 are presented. No new matter has been added.

004090-67403550

Applicant submits that all of the claims are now in condition for examination, which action is requested. Filed herewith is a check in payment of the excess claims fees required by the above amendments. Please apply any other charges or credits to Deposit Account No. 06-1050.

Respectfully submitted,

Date:\_\_\_\_\_

Chris T. Mizumoto  
Reg. No. 42,899

Fish & Richardson P.C.  
45 Rockefeller Plaza, Suite 2800  
New York, NY 10111  
Telephone: (212) 765-5070  
Facsimile: (212) 258-2291

30019137 doc

Table 1. Demographic characteristics of the study population	
<b>Age (years)</b>	
18-24	10 (10.0%)
25-34	15 (15.0%)
35-44	20 (20.0%)
45-54	25 (25.0%)
55-64	30 (30.0%)
65-74	35 (35.0%)
75-84	40 (40.0%)
85-94	45 (45.0%)
95-104	50 (50.0%)
105-114	55 (55.0%)
115-124	60 (60.0%)
125-134	65 (65.0%)
135-144	70 (70.0%)
145-154	75 (75.0%)
155-164	80 (80.0%)
165-174	85 (85.0%)
175-184	90 (90.0%)
185-194	95 (95.0%)
195-204	100 (100.0%)
<b>Gender</b>	
Male	100 (100.0%)
Female	0 (0.0%)
<b>Ethnicity</b>	
White	100 (100.0%)
Black	0 (0.0%)
Hispanic	0 (0.0%)
Asian	0 (0.0%)
Other	0 (0.0%)
<b>Marital status</b>	
Married	100 (100.0%)
Single	0 (0.0%)
Divorced	0 (0.0%)
Widowed	0 (0.0%)
<b>Education level</b>	
High school or less	100 (100.0%)
Some college	0 (0.0%)
Bachelor's degree	0 (0.0%)
Master's degree	0 (0.0%)
PhD	0 (0.0%)
<b>Occupation</b>	
Unemployed	100 (100.0%)
Employed	0 (0.0%)
<b>Health status</b>	
Good	100 (100.0%)
Fair	0 (0.0%)
Poor	0 (0.0%)
<b>Smoking status</b>	
Smoker	100 (100.0%)
Non-smoker	0 (0.0%)
<b>Alcohol consumption</b>	
Drinker	100 (100.0%)
Non-drinker	0 (0.0%)
<b>Exercise frequency</b>	
Regular	100 (100.0%)
Occasional	0 (0.0%)
Never	0 (0.0%)
<b>Family size</b>	
1-2	100 (100.0%)
3-4	0 (0.0%)
5-6	0 (0.0%)
7-8	0 (0.0%)
9-10	0 (0.0%)
11-12	0 (0.0%)
13-14	0 (0.0%)
15-16	0 (0.0%)
17-18	0 (0.0%)
19-20	0 (0.0%)
21-22	0 (0.0%)
23-24	0 (0.0%)
25-26	0 (0.0%)
27-28	0 (0.0%)
29-30	0 (0.0%)
31-32	0 (0.0%)
33-34	0 (0.0%)
35-36	0 (0.0%)
37-38	0 (0.0%)
39-40	0 (0.0%)
41-42	0 (0.0%)
43-44	0 (0.0%)
45-46	0 (0.0%)
47-48	0 (0.0%)
49-50	0 (0.0%)
51-52	0 (0.0%)
53-54	0 (0.0%)
55-56	0 (0.0%)
57-58	0 (0.0%)
59-60	0 (0.0%)
61-62	0 (0.0%)
63-64	0 (0.0%)
65-66	0 (0.0%)
67-68	0 (0.0%)
69-70	0 (0.0%)
71-72	0 (0.0%)
73-74	0 (0.0%)
75-76	0 (0.0%)
77-78	0 (0.0%)
79-80	0 (0.0%)
81-82	0 (0.0%)
83-84	0 (0.0%)
85-86	0 (0.0%)
87-88	0 (0.0%)
89-90	0 (0.0%)
91-92	0 (0.0%)
93-94	0 (0.0%)
95-96	0 (0.0%)
97-98	0 (0.0%)
99-100	0 (0.0%)
101-102	0 (0.0%)
103-104	0 (0.0%)
105-106	0 (0.0%)
107-108	0 (0.0%)
109-110	0 (0.0%)
111-112	0 (0.0%)
113-114	0 (0.0%)
115-116	0 (0.0%)
117-118	0 (0.0%)
119-120	0 (0.0%)
121-122	0 (0.0%)
123-124	0 (0.0%)
125-126	0 (0.0%)
127-128	0 (0.0%)
129-130	0 (0.0%)
131-132	0 (0.0%)
133-134	0 (0.0%)
135-136	0 (0.0%)
137-138	0 (0.0%)
139-140	0 (0.0%)
141-142	0 (0.0%)
143-144	0 (0.0%)
145-146	0 (0.0%)
147-148	0 (0.0%)
149-150	0 (0.0%)
151-152	0 (0.0%)
153-154	0 (0.0%)
1	

004090" 6T45B550

**APPLICATION**  
**FOR**  
**UNITED STATES LETTERS PATENT**

**TITLE:** **DIAGNOSTIC AGENTS FOR PANCREATIC EXOCRINE  
FUNCTION**

**APPLICANT:** **TADASHI KOHNO, ISABURO HOSOI, JUNKO OHSHIMA,  
KUNIHICO SHIBATA & ASUKA ITO**

"EXPRESS MAIL" Mailing Label Number EI372100430US

Date of Deposit September 23, 1999

I hereby certify under 37 CFR 1.10 that this correspondence is being deposited with the United States Postal Service as "Express Mail Post Office To Addressee" with sufficient postage on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231.

Francisco Robles

Francisco Robles



## SPECIFICATION

### DIAGNOSTIC AGENTS FOR PANCREATIC EXOCRINE FUNCTION

#### BACKGROUND OF THE INVENTION

##### Field of the Invention

The present invention relates to novel compounds useful as diagnostic agents for pancreatic exocrine function and their use.

##### Background of the Invention

"Pancreatic exocrine function tests" are useful for the diagnosis of pancreatic diseases such as chronic and acute pancreatitis and pancreatic cancer. They are also useful to assess the condition and prognosis of patients and to control the medication: The general descriptions are found in Arvanitakis and Cooke, Gastroenterology, 74:932 (1978); Niederau and Grendell, Gastroenterology, 88:1973 (1985); Goldberg, Bull. Mol. Biol. Med., 15:1 (1990); Lankisch, Int. J. Pancreatology, 14:9 (1993); Bank and Chow, Gastroenterologist, 2:224 (1994); and Steer et al., New Eng. J. Med., 332:1482 (1995).

At present, "Gold standard" of the pancreatic exocrine function test involves inserting a tube through the mouth to the duodenum to collect the duodenal juice. Now, the secretin test is generally utilized wherein secretin is intravenously administered to stimulate the secretion of the pancreatic juice prior to the collection. This method is highly accurate since the amount and components of the pancreatic juice are directly analyzed. However, this method can not be used repeatedly or used for screening because of the very strong stress caused on the subjects. It is available at only a relatively small number of medical centers having highly skilled physicians. Further, since this method requires fluoroscopic tube placement during the collection of the duodenal juice, there is the problem of X

ray exposure.

On the other hand, a test for quantifying pancreatic exocrine enzymes from the pancreas into the blood is clinically employed for screening pancreatic diseases (The Merck Manual 16th edition). However, the increase of the pancreatic exocrine enzymes in the blood is only observed at the initial stage of acute pancreatitis or at the recrudescence stage of chronic pancreatitis and does not always reflect the ability of the pancreas to secrete pancreatic exocrine enzymes. Further, the increase of pancreatic exocrine enzymes in the blood may sometimes not be detected due to serum turbidity in pancreatitis accompanied by hyperlipemia.

Accordingly, simple methods which require no insertion of a tube are utilized for repetition and screening tests. One of them is the pancreolauryl test (PLT) wherein a synthetic substrate FDL (fluorescein dilaurylate, dilaurylfluorescein) for cholesterol ester hydrolase, esterase, secreted from the pancreas is orally administered and urine is accumulated for 10 hours followed by measuring the amount of a degradation product fluorescein excreted into the urine: US Patent No. 3,917,812; Barry et al., Lancet (1982) Oct. 2, p. 742; Scharpe and Iliano, Clin. Chem., 33:5 (1987). However, this method requires a long time to carry out the test and therefore can not often be performed on outpatients and is not suitable in physical examinations.

Under these circumstances, there is a need for the development of a simple method for testing the pancreatic exocrine function which imparts low stress on subjects and gives accurate results soon.

On the one hand, the  $^{13}\text{C}$ -breath test wherein a  $^{13}\text{C}$ -labeled starch is administered has been recently considered to be employed in the test for the pancreatic exocrine function: Hiele et al., Gastroenterology,

96:503 (1989); Dewit et al., Pediatric Res., 32:45 (1992); and Z. Gastroenterol., 35:187 (1997). In the enteric tract, starch is degraded efficiently to glucose by the cleavage at any internal  $\alpha$ -1,4 glucoside linkage with  $\alpha$ -amylase secreted from the pancreas and by the action of enzymes such as  $\alpha$ -glucosidase (maltase) of mucosal epithelial cells of the small intestine and absorbed: Essentials of Human Metabolism, 2nd ed., W.C. McMurray, Harper & Row Publishers, NY. The  $^{13}\text{C}$ -breath test wherein a  $^{13}\text{C}$ -labeled starch is administered utilizes the phenomenon that after the  $^{13}\text{C}$ -labeled starch is degraded in the digestive tract, it is absorbed and decarboxylated by metabolic action in the body to generate  $^{13}\text{CO}_2$ , which is excreted into the breath, and it is a safe and simple method. However, any  $^{13}\text{C}$ -labeled oligosaccharide or polysaccharide other than  $^{13}\text{C}$ -labeled starch has not yet been studied.

Since there is an  $\alpha$ -glucosidase (maltase) in mucosal epithelial cells of the small intestine, which cleaves a non-reducing terminal  $\alpha$ -1,4- glucoside linkage (Enzyme Handbook, Springer-Verlag, Berlin), starch is degraded into glucose sequentially from the non-reducing terminal and absorbed only by the action of enzymes such as the  $\alpha$ -glucosidase (maltase), even without the action of  $\alpha$ -amylase. Thus, starch is subject to the action of a non-pancreatic-exocrine enzyme  $\alpha$ -glucosidase (maltase) of mucosal epithelial cells of the small intestine and therefore the  $^{13}\text{C}$ -labeled starch breath test does not reflect the pancreatic exocrine function only. Accordingly, it would be more preferred if a substrate compound specific for  $\alpha$ -amylase in the digestive tract is selected.

Accordingly, an object of the present invention is to

provide a diagnostic agent for pancreatic exocrine function which leads to a test for the pancreatic exocrine function imparting low stress on subjects and yields the results in a short period of time.

It is another object of the present invention to provide a diagnostic agent for pancreatic exocrine function which is specific to the  $\alpha$ -amylase secretion ability.

It is a further object of the present invention to provide a novel compound usable in the pancreatic exocrine function test.

#### SUMMARY OF THE INVENTION

The present inventors have found that the pancreatic exocrine function test can be carried out by orally administering a  $^{13}\text{C}$ -labeled oligosaccharide or a  $^{13}\text{C}$ -labeled inclusion complex or a  $^{13}\text{C}$ -labeled fluorescein ester compound to a rat with chronic pancreatitis and measuring the  $^{13}\text{C}$  concentration in the exhaled  $\text{CO}_2$  after administration. Thus, the present invention has been completed.

Accordingly, the present invention provides a  $^{13}\text{C}$ -labeled oligosaccharide or polysaccharide or a salt thereof or a derivative thereof or a  $^{13}\text{C}$ -labeled inclusion complex other than  $\text{U-}^{13}\text{C}$ -maltose,  $^{13}\text{C}$ -starch,  $1\text{-}^{13}\text{C}$ -maltotetraose and  $1\text{-}^{13}\text{C}$ -amylose.

Also, the present invention provides a diagnostic agent for pancreatic exocrine function comprising a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled oligosaccharide or polysaccharide or a salt or a derivative thereof, or a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled inclusion complex or a salt thereof other than  $^{13}\text{C}$ -starch.

Moreover, the present invention provides a diagnostic agent for pancreatic exocrine function comprising a compound which is degraded with  $\alpha$ -amylase

but not degraded with  $\alpha$ -glucosidase.

Furthermore, the present invention provides a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled fluorescein ester compound or a salt thereof.

Also, the present invention provides a diagnostic agent for pancreatic exocrine function comprising a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled fluorescein ester compound or a pharmaceutically acceptable salt thereof.

The subject matters of the present invention are as follows.

(1) A  $^{13}\text{C}$ -labeled oligosaccharide or polysaccharide or a salt thereof excluding U- $^{13}\text{C}$ -maltose,  $^{13}\text{C}$ -starch, 1- $^{13}\text{C}$ -maltotetraose and 1- $^{13}\text{C}$ -amylose.

(2) The  $^{13}\text{C}$ -labeled oligosaccharide or polysaccharide or salt thereof according to (1), which is hydrolyzed with  $\alpha$ -amylase.

(3) The  $^{13}\text{C}$ -labeled oligosaccharide or polysaccharide or salt thereof according to (2), which is not hydrolyzed with  $\alpha$ -glucosidase.

(4) The  $^{13}\text{C}$ -labeled oligosaccharide or polysaccharide or salt thereof according to any one of (1) to (3), wherein at least one sugar molecule constituting the oligosaccharide or polysaccharide is  $^{13}\text{C}$ -labeled.

(5) The  $^{13}\text{C}$ -labeled oligosaccharide or polysaccharide or salt thereof according to any one of (1) to (3), wherein at least one sugar molecule constituting the oligosaccharide or polysaccharide is modified with at least one  $^{13}\text{C}$ -labeled modifying group.

(6) The  $^{13}\text{C}$ -labeled oligosaccharide or polysaccharide or salt thereof according to any one of (1) to (5), which is a linear or branched oligosaccharide or polysaccharide.

(7) The  $^{13}\text{C}$ -labeled oligosaccharide or

polysaccharide or salt thereof according to any one of (1) to (5), which is a cyclic oligosaccharide or polysaccharide.

(8) The  $^{13}\text{C}$ -labeled oligosaccharide or polysaccharide or salt thereof according to (6), which is modified at the non-reducing terminal.

(9) The  $^{13}\text{C}$ -labeled oligosaccharide or polysaccharide or salt thereof according to (1), which is a  $^{13}\text{C}$ -cyclodextrin or  $\beta$ -galactosyl- $^{13}\text{C}$ -maltooligosaccharide.

(10) A derivative of the  $^{13}\text{C}$ -labeled oligosaccharide or polysaccharide or salt thereof according to any one of (1) to (9).

(11) A  $^{13}\text{C}$ -labeled inclusion complex or a salt thereof, which comprises a cyclodextrin or a modified derivative thereof as a host molecule.

(12) The inclusion complex or salt thereof according to (11), wherein the host molecule is  $^{13}\text{C}$ -labeled.

(13) The inclusion complex or salt thereof according to (11), wherein the guest molecule is  $^{13}\text{C}$ -labeled.

(14) The inclusion complex or salt thereof according to any one of (11) to (13), wherein the guest molecule is selected from the group consisting of oligosaccharides, amino acids, peptides, organic acids, fatty acids, fatty acid glycerides, vitamins, catechins, carotinoids, flavonoids and cholesterol.

(15) The inclusion complex or salt thereof according to (14), wherein the guest molecule is selected from the group consisting of  $^{13}\text{C}$ -phenylalanine, benzoylphenylalanyl- $^{13}\text{C}$ -leucine and benzoylphenylalanyl- $^{13}\text{C}$ -leucine methyl ester.

(16) A diagnostic agent for pancreatic exocrine



ability of the pancreas to secrete  $\alpha$ -amylase and at least one pancreatic exocrine enzyme other than  $\alpha$ -amylase.

(25) A  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled fluorescein ester compound or a salt thereof.

(26) The compound or salt thereof according to (25), which is a compound or salt thereof resulting from a reaction of a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled acid with fluorescein at both or either of the two hydroxyl groups at 3' and 6' positions to form an ester linkage.

(27) The compound or salt thereof according to (26), wherein the acid is a carboxylic acid.

(28) The compound or salt thereof according to (27), wherein the carboxylic acid is a fatty acid.

(29) The compound or salt thereof according to (28), wherein the fatty acid has 2 to 16 carbon atoms.

(30) The compound or salt thereof according to (29), wherein the fatty acid is selected from the group consisting of lauric, acetic and octanoic acids.

(31) The  $^{13}\text{C}$ -labeled fluorescein ester compound or salt thereof according to (25), which is selected from the group consisting of the following compounds:

- (a)  $^{13}\text{C}$ -dilaurylfluorescein;
- (b)  $^{13}\text{C}$ -diacetylfluorescein; and
- (c)  $^{13}\text{C}$ -dioctanoylfluorescein.

(32) A diagnostic agent for pancreatic exocrine function comprising a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled fluorescein ester compound or a pharmaceutically acceptable salt thereof.

(33) The diagnostic agent for pancreatic exocrine function according to (32), comprising a compound or salt thereof resulting from a reaction of a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled acid with fluorescein at both or either of the two hydroxyl groups at 3'



and 6' positions to form an ester linkage.

(34) The diagnostic agent for pancreatic exocrine function according to (33), wherein the acid is a carboxylic acid.

(35) The diagnostic agent for pancreatic exocrine function according to (34), wherein the carboxylic acid is a fatty acid.

(36) The diagnostic agent for pancreatic exocrine function according to (35), wherein the fatty acid has 2 to 16 carbon atoms.

(37) The diagnostic agent for pancreatic exocrine function according to (32), which is selected from the group consisting of the following compounds:

(a)  $^{13}\text{C}$ -dilaurylfluorescein;

(b)  $^{13}\text{C}$ -diacetylfluorescein; and

(c)  $^{13}\text{C}$ -dioctanoylfluorescein.

(38) The diagnostic agent for pancreatic exocrine function according to any one of (32) to (37), wherein the  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled fluorescein ester compound or pharmaceutically acceptable salt thereof is subjected to the action of the pancreatic exocrine cholesterol ester hydrolase and pancreatic exocrine esterase and decarboxylated to generate  $^{13}\text{CO}_2$  or  $^{14}\text{CO}_2$ .

(39) A  $^{13}\text{C}$ -labelled oligosaccharide or polysaccharide or salt thereof according to any one of claims (1) to (10) for use in a method of diagnosis.

(40) An inclusion complex according to any of claims (11) to 15 for use in a method of diagnosis.

(41) A  $^{13}\text{C}$  or  $^{14}\text{C}$ -labelled fluorescein ester compound or salt thereof according to any one of claims (25) to (31) for use in a method of diagnosis.

(42) A compound according to any one of claims (39) to (41) wherein the diagnosis is the diagnosis of pancreatic exocrine function.



The term "branched oligosaccharide or polysaccharide" refers to an oligosaccharide or polysaccharide having a branched structure such as amylopectin and glycogen.

The term "cyclic oligosaccharide or polysaccharide" refers to an oligosaccharide or polysaccharide having a cyclic structure such as cyclodextrin.

The term "non-reducing terminal" refers to the terminal at the side at which the carbon atom at position 1 of a sugar residue is involved in binding of the sugar chain.

This specification includes part or all of the contents as disclosed in the specifications and/or drawings of Japanese Patent Application Nos. 10-2712152, 10-271253, 11-261979 and 11-263300 which are priority documents of the present application.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 shows a  $^{13}\text{C}$ -NMR spectrum of  $^{13}\text{C}$ -cyclodextrin.

Fig. 2 shows the time course of degree of increase of the  $^{13}\text{C}$  concentration in the exhaled  $\text{CO}_2$  ( $\Delta^{13}\text{C}$  (%)) after administration of  $^{13}\text{C}$ -labeled cyclodextrin. At 0 minute,  $^{13}\text{C}$ -labeled cyclodextrin was orally administered (75 mg/kg) to chronic pancreatitis rats (n=2, ●) and control rats (n=4, ○). The bar indicates SD.

Fig. 3 shows a  $^{13}\text{C}$ -NMR spectrum of galactosyl  $^{13}\text{C}$ -maltohexaose.

Fig. 4 shows the time course of degree of increase of the  $^{13}\text{C}$  concentration in the exhaled  $\text{CO}_2$  ( $\Delta^{13}\text{C}$  (%)) after administration of  $^{13}\text{C}$ -galactosyl maltohexaose. At 0 minutes,  $^{13}\text{C}$ -galactosyl maltohexaose was orally administered (75 mg/kg) to chronic pancreatitis rats (n=3, ●) and control rats (n=4, ○). The bar indicates SD.

Fig. 5 shows the time course of degree of increase of the  $^{13}\text{C}$  concentration in the exhaled  $\text{CO}_2$  ( $\Delta^{13}\text{C}$  (%)) after administration of  $^{13}\text{C}$ -labeled phenylalanine/hydroxypropyl- $\beta$ -cyclodextrin ( $1\text{-}^{13}\text{C-Phe/hp-}\beta\text{-CD}$ ) inclusion complex. At 0 minutes,  $1\text{-}^{13}\text{C-Phe/hp-}\beta\text{-CD}$  inclusion complex was orally administered (59.76 mg/kg of  $1\text{-}^{13}\text{C-Phe}$ ) to chronic pancreatitis rats ( $n=4$ , ●) and control rats ( $n=4$ , ○). The bar indicates SD.

Fig. 6 is a powder X-ray diffraction spectrum of benzoylphenylalanyl [ $1\text{-}^{13}\text{C}$ ]-leucine/ $\gamma$ -cyclodextrin (Bz-Phe- ( $^{13}\text{C-Leu}$ )/ $\gamma$ -CD) inclusion complex.

Fig. 7 shows the time course of degree of increase of the  $^{13}\text{C}$  concentration in the exhaled  $\text{CO}_2$  ( $\Delta^{13}\text{C}$  (%)) after administration of benzoylphenylalanyl [ $1\text{-}^{13}\text{C}$ ]-leucine/ $\gamma$ -cyclodextrin (Bz-Phe- ( $^{13}\text{C-Leu}$ )/ $\gamma$ -CD) inclusion complex. At 0 minute, Bz-Phe- ( $^{13}\text{C-Leu}$ )/ $\gamma$ -CD was orally administered (100 mg/kg of Bz-Phe- ( $^{13}\text{C-Leu}$ )) to chronic pancreatitis rats ( $n=2$ , ●) and control rats ( $n=2$ , ○). The bar indicates SD.

Fig. 8 is a powder X-ray diffraction spectrum of benzoylphenylalanyl [ $1\text{-}^{13}\text{C}$ ]-leucine methyl ester/ $\gamma$ -cyclodextrin (Bz-Phe- ( $^{13}\text{C-Leu}$ )Me/ $\gamma$ -CD) inclusion complex.

Fig. 9 shows the time course of degree of increase of the  $^{13}\text{C}$  concentration in the exhaled  $\text{CO}_2$  ( $\Delta^{13}\text{C}$  (%)) after administration of benzoylphenylalanyl [ $1\text{-}^{13}\text{C}$ ]-leucine methyl ester/ $\gamma$ -cyclodextrin (Bz-Phe- ( $^{13}\text{C-Leu}$ )Me/ $\gamma$ -CD) inclusion complex. At 0 minute, Bz-Phe- ( $^{13}\text{C-Leu}$ )Me/ $\gamma$ -CD was orally administered (70 mg/kg of Bz-Phe- ( $^{13}\text{C-Leu}$ )Me) to chronic pancreatitis rats ( $n=4$ , ●) and control rats ( $n=4$ , ○). The bar indicates SD.

Fig. 10 shows the time course of degree of increase of the

$^{13}\text{C}$  concentration in the exhaled  $\text{CO}_2$  ( $\Delta^{13}\text{C}$  (%)) after administration of  $^{13}\text{C}$ -dilaurylfluorescein ( $^{13}\text{C}$ -FDL). At 0 minute,  $^{13}\text{C}$ -FDL was orally administered (160 mg/kg) to chronic pancreatitis rats (●, n=3) and control rats (□, n=3). Bars represent SD.

Fig. 11 shows the time course of degree of increase of the  $^{13}\text{C}$  concentration in the exhaled  $\text{CO}_2$  ( $\Delta^{13}\text{C}$  (%)) after administration of  $^{13}\text{C}$ -dioctanoylfluorescein ( $^{13}\text{C}$ -FDO). At 0 minute,  $^{13}\text{C}$ -FDO was orally administered (200 mg/kg) to chronic pancreatitis rats ( $\bullet$ , n=4) and control rats ( $\square$ , n=4). Bars represent SD.

## DESCRIPTION OF THE INVENTION

Hereinafter, the present invention will be described in detail.

The present invention encompasses  $^{13}\text{C}$ -labeled oligosaccharide or polysaccharides or salts thereof, or derivatives thereof or  $^{13}\text{C}$ -labeled inclusion complex other than U- $^{13}\text{C}$ -maltose,  $^{13}\text{C}$ -starch, 1- $^{13}\text{C}$ -maltotetraose and 1- $^{13}\text{C}$ -amylose. The diagnostic agent for pancreatic exocrine function of the present invention comprises a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled oligosaccharide or polysaccharide or a salt thereof a derivative thereof other than  $^{13}\text{C}$ -starch, or a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled inclusion complex or a salt thereof. Preferably, the  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled oligosaccharide or polysaccharides or salts thereof, or derivatives thereof are modified at the non-reducing terminals or have cyclic structure. These  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled compounds may be pharmaceutically acceptable.

The term "<sup>13</sup>C- or <sup>14</sup>C-labeled" herein means that any carbon atom in the compound is replaced with a <sup>13</sup>C or <sup>14</sup>C atom, resulting in a higher abundance ratio of <sup>13</sup>C or <sup>14</sup>C

than the naturally occurring abundance ratio irrespective of the preparation method thereof.

The "oligosaccharide or polysaccharides modified at their non-reducing terminals" refers to oligosaccharide or polysaccharides in which the carbon atom on their non-reducing terminal glucosyl group is modified. Modifying groups in addition to glucopyranose include modifying groups comprising oligosaccharides such as galactosyl and digalactosyl groups, alkyl groups, alkoxyl groups such as methoxy and benzyloxy groups, carbamoyl group, pyridylamino group, and ethylidene and benzylidene groups which are attached to two carbon atoms on the non-reducing terminal glucosyl group.

Examples of the oligosaccharide or polysaccharide or salt thereof, or derivative thereof include paranitrophenyl-6-o-benzyl maltopentaoside, 4-nitrophenyl maltohexaoside 4,6-ethylidene glucoside, 4,6-benzylidene  $\alpha$ -o-4-nitrophenyl-maltopentaoside, o-6-deoxypyridylamino- $\alpha$ -maltopentaoside and paranitrophenyl-4,6-di-o-(N-ethyl)4-carbamoyl maltopentaoside and so on.

These compounds can be prepared by methods described in the following literature:

Carbohydrate Research, 176 (1988), 107-115;

Japanese Patent Application Laying Open No. 3-91496;

US Patent No. 4,649,108;

Journal of Chromatography, 336 (1984), 368-373; and

Japanese Patent Application Laying Open No. 8-291

In the preparation, a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled maltooligosaccharide (which may be commercially available or be prepared by a known method) can be used as a starting material to give a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled one.

For example, galactosyl oligosaccharide, which is an oligosaccharide modified at the non-reducing terminal, may be obtained by adding lactose to an oligosaccharide

to bring lactase into action thereon, and collecting from the reaction mixture by column chromatography (Japanese Patent Application Laying Open Nos. 4-209277, 8-196289 and 10-316697). In the preparation, a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled oligosaccharide can be used to give a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled galactosyl oligosaccharide. When a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -galactosyl labeled lactose is used, an galactosyloligosaccharide in which the galactosyl group is  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled is obtained.

"Cyclodextrins" herein means  $\alpha$ -1,4 linked cyclic oligosaccharides composed of glucopyranose units. The number of glucoses forming the ring is not limited at all and  $\alpha$ -,  $\beta$ - and  $\gamma$ -cyclodextrins having 6 to 8 glucoses are commercially available. The cyclodextrin may have a branch. The hydroxyl group(s) in cyclodextrin may be modified; cyclodextrins having such a modifying group are included in the "modified derivatives of cyclodextrin". Illustratively, the modification includes alkylation such as methylation, hydroxyalkylation such as hydroxypropylation, esterification such as acetylation or succinylation, glucosylation, carboxymethyl etherification, phosphoric esterification, sulfobutyl etherification, and carboxymethylation. These modified derivatives are described in Loftsson et al., J. Pharmaceu. Sci., 85:1017 (1996); Stella et al., Pharmaceutical Res., 14:556 (1997); and "Cyclodextrins" supervised by Fujio Toda, published by Sangyo Tosho.

For example, a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled cyclodextrin may be obtained by bringing a cyclomaltodextrin glucanotransferase into action on a naturally occurring  $^{13}\text{C}$ -enriched starch derived from C4 plants such as corn or a commercially available  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled starch and collecting from the reaction mixture by column chromatography.





are oligosaccharides, peptides, fatty acid glycerides and modified derivatives thereof.

The  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled cyclodextrin inclusion complex may be one in which any carbon atom in the included compound is replaced with a  $^{13}\text{C}$  or  $^{14}\text{C}$  atom, resulting in a higher abundance ratio of  $^{13}\text{C}$  or  $^{14}\text{C}$  in the cyclodextrin inclusion complex than the naturally occurring abundance ratio. However, the cyclodextrin or modified derivative thereof per se may be  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled. Methods for the preparation thereof are not limited.

Examples of cyclodextrin inclusion complexes reported heretofore include the following (host molecules are not limited to ones indicated in the complexes):

Phenylalanine/ $\beta$ -cyclodextrin inclusion complex

Tryptophan/ $\alpha$ -cyclodextrin inclusion complex

Histidine/ $\alpha$ -cyclodextrin inclusion complex

Cinnalidine/ $\beta$ -cyclodextrin inclusion complex

Ferulic acid/ $\beta$ -cyclodextrin inclusion complex

Phenylalanyl leucine/ $\beta$ -cyclodextrin inclusion complex


Other guest molecules include phenol, hydroxy benzoic acid, benzaldehyde, methyl sulfoxide, benzoic acid, aniline, amino benzoic acid, methyl benzoic acid, nitrophenol, pyridine, acetic acid, alcohols (e.g., ethanol, propanol, butanediol, etc.), prostandin, benexate hydrochloride, nitroglycerin, limaprost and the like, which are described in "Cyclodextrin" edited by Fujio Toda, Sangyo Tosho. The cyclodextrin inclusion complex can be obtained by mixing purified water with a mixture of a host compound and a guest compound at a ratio equal to that of the host to the guest in the inclusion complex (e.g., at an equivalent molar ratio in the case where the ratio of the host to the guest is 1:1),

stirring the resulting mixture for about 12 hours, and subjecting it to a spray dry treatment.

The  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled oligosaccharide or polysaccharide or cyclodextrin inclusion complex may be also obtained in the form of a salt. Such salts include those with inorganic acids such as hydrochloric, sulfuric and phosphoric acids; those with organic acids such as formic, acetic, propionic, glycolic, succinic, malic, tartaric, citric and trifluoroacetic acids; those with alkali metals such as sodium and potassium; those with alkaline earth metals such as calcium; and those with ammonium or organic amines such as ethanolamine, triethylamine and dicyclohexylamine.

The present invention also encompasses  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled fluorescein ester compounds or salts thereof.

Fluorescein (CA Name: 3',6'-dihydroxyspiro[isobenzofuran-1(3H),9'-[9H]xanthen-3-one) is represented by the following structural formula:



The "fluorescein ester compound" refers to a compound resulting from a reaction of an acid with fluorescein at the hydroxyl group(s) to form an ester linkage. Fluorescein and the acid may be modified.

The " $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled" means that at least one carbon atom in the fluorescein ester compound is replaced with a  $^{13}\text{C}$ - or  $^{14}\text{C}$  atom, resulting in a higher abundance ratio of the  $^{13}\text{C}$ - or  $^{14}\text{C}$  atom than the naturally occurring abundance ratio.



function according to the present invention may be formulated from the  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled compound alone or in combination with an excipient or carrier into an oral preparation such as a tablet, capsule, powder, granule or liquid. The excipient or carrier may be any pharmaceutically acceptable one ordinarily used in this field and its nature and composition may be appropriately chosen. For example, water may be used as a liquid carrier. Solid carriers include cellulose derivatives such as hydroxypropyl cellulose, and organic acid salts such as magnesium stearate. Also, freeze-dried preparations may be used.

The  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled compound is contained in the preparation in variable amounts depending on the nature of the preparation, but generally in an amount of 1 to 100% by weight, preferably 50 to 100% by weight. In a capsule, tablet, granule or powder preparation, the  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled compound is contained in the preparation in an amount of about 10 to 100% by weight, preferably 50 to 100% by weight, the balance being a carrier.

The dose of the diagnostic agent for pancreatic exocrine function according to the present invention should be sufficient to confirm an increase of  $^{13}\text{CO}_2$  or  $^{14}\text{CO}_2$  in the breath after the administration. It will vary depending upon the age and body weight of a subject and the purpose of the test. For example, the unit dose may be 1 to 2000 mg/kg of body weight for an adult.

The test using the agent for pancreatic exocrine function according to the present invention is carried out with administering to a subject the  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled compound. A test is possible, in which the concentration of a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled compound is measured in serum, urine or stool after the administration, however, a breath test is desirable in

which an increase in  $^{13}\text{C}$  or  $^{14}\text{C}$  concentration is measured in the exhaled  $\text{CO}_2$  after the administration. When the  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled compound is administered to a subject, a test meal or the like may be pre-ingested by the subject to induce secretion of pancreatic enzymes. The  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled compound may be administered together with the test meal or the like. Also, a plurality of the  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled compound may be combined for use. Concretely, in the case of  $^{13}\text{C}$ , the  $^{13}\text{C}$  concentration is determined in the exhaled  $\text{CO}_2$  after the administration, then the pancreatic exocrine function is diagnosed from either the data of the degree of increase ( $\Delta^{13}\text{C}$  (%)) of the  $^{13}\text{C}$  concentration in the exhaled  $\text{CO}_2$  at predetermined times (e.g., 5, 10 and 15 minutes) after the administration, or the data integrated or associated with the time course (onset slope, change in slope, peak time, etc.) in the degree of increase ( $\Delta^{13}\text{C}$  (%)) of the  $^{13}\text{C}$  concentration in the exhaled  $\text{CO}_2$  during a predetermined period after the administration. In the case of  $^{14}\text{C}$ , the  $^{14}\text{C}$  concentration, i.e., radioactivity, is determined in the exhaled  $\text{CO}_2$  after the administration; and the pancreatic exocrine function is diagnosed from either the data of the quantity of radioactivity in the exhaled  $\text{CO}_2$  at predetermined times (e.g., 5, 10 and 15 minutes) after the administration, or the data integrated or associated with the time course (onset slope, change in slope, peak time, etc.) in the rate increase of radioactivity in the exhaled  $\text{CO}_2$  during a predetermined period after the administration.

These test methods utilize the phenomenon that when the  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled compound is administered to a subject, the compound is absorbed through the digestive tract after it is degraded by the action of the pancreatic exocrine  $\alpha$ -amylase and/or esterase, and



## EXAMPLES

### Example 1: Preparation of $^{13}\text{C}$ -labeled cyclodextrin

$^{13}\text{C}$ -labeled starch (Chlorella Industry, Algal Starch (water-soluble), Lot No. 8031,S, U- $^{13}\text{C}$ :98.6 atom %, Starch Content: 93.5%) was dissolved in 50 mM acetate buffer (pH 5.4) at a concentration of 5% (w/v) and 100 Units of cyclomaltodextrin glucanotransferase (Hayashibara) was added thereto and reacted at 40°C for 24 hours. After the enzyme was inactivated with the treatment at 100°C for 15 minutes, glucoamylase was added and reacted at 40°C for 1 hour to decompose components other than  $^{13}\text{C}$ -labeled cyclodextrin into glucose. After the reaction was completed, the reaction mixture was treated at 100°C for 15 minutes to inactivate the glucoamylase.

The solution was applied to a carbon column (2.5 cm x 25 cm) and the column was washed with 500 ml of water. A 15% ethanol solution and a 40% ethanol solution were sequentially applied to recover the  $^{13}\text{C}$ -labeled cyclodextrin in the 40% ethanol solution eluted fractions. The resulting  $^{13}\text{C}$ -labeled cyclodextrin solution was distilled to remove the solvent, dissolved in a small amount of water and freeze-dried.

The product was mixed with para-nitrophenol to confirm to be cyclodextrin from an increase in absorbance at a wave length of 450 nm.

The  $^{13}\text{C}$ -labeled position was confirmed by  $^{13}\text{C}$ -NMR (Fig.1).

$^{13}\text{C}$ -NMR (DMSO-d<sub>6</sub>, 300 MHz)

38.5-40.2 ppm	Dimethyl sulfoxide
59.5-60.1 ppm	Position 6 of glucose residue
71.4-73.1 ppm	Positions 3, 4 and 5 of glucose residue
80.9-82.5 ppm	Position 2 of glucose residue

101.5-102.1 ppm Position 1 of glucose residue

HPLC analysis (Shodex Asahipak GS-220 HQ) of the resulting  $^{13}\text{C}$ -labeled cyclodextrin revealed that it was a mixture of 45% of  $\alpha$ -cyclodextrin, 45% of  $\beta$ -cyclodextrin and 10% of  $\gamma$ -cyclodextrin.

#### Example 2: $^{13}\text{C}$ -labeled cyclodextrin breath test

$^{13}\text{C}$ -labeled cyclodextrin breath test was carried out wherein  $^{13}\text{C}$ -labeled cyclodextrin prepared in Example 1 was orally administered to chronic pancreatitis and control rats and the time course of the  $^{13}\text{C}$  concentration in the exhaled  $\text{CO}_2$  after the administration was measured.

According to Mundlos et al. (Mundlos et al., Pancreas, 1:29 (1986)), the chronic pancreatitis rats were prepared by injecting oleic acid into the pancreatic duct of Wistar male rats of 5 weeks old and kept for 3 weeks. Rats in which midline incision was made on the abdomen were used as the control.

The chronic pancreatitis and control rats of 8 weeks old, which fasted overnight, were fixed without anesthesia in a rat holder for a microwave irradiation apparatus. The breath was collected at a rate of about 100 to 300 ml/min using a stroke pump (Variable Stroke Pump VS-500, Shibata Kagaku Kogyo) and introduced directly to a flow cell of a  $^{13}\text{CO}_2$  analyzer EX-130S (Nihon Bunko) to measure  $^{13}\text{C}$  atom% and the carbon dioxide gas concentration continuously. A Perma Pure drier (MD-050-12P, Perma Pure Inc.) was set between the rat holder and the stroke pump to remove out water vapor in the breath. After the  $\text{CO}_2$  concentration was stabilized, the rat was once removed out of the rat holder and an aqueous solution of  $^{13}\text{C}$ -labeled cyclodextrin was administered (75 mg/kg, 5 ml/kg) into the stomach using an oral sonde.

$\Delta^{13}\text{C}$  (‰) was calculated from the  $^{13}\text{C}$  concentration



in the exhaled CO<sub>2</sub> at each time point (<sup>13</sup>C tmin) and the <sup>13</sup>C concentration in standard CO<sub>2</sub> (<sup>13</sup>C std) according to the following equation:

$$\Delta^{13}\text{C} (\%) = [({}^{13}\text{C tmin} - {}^{13}\text{C 0min}) / {}^{13}\text{C std}] \times 1000$$

In both the control and chronic pancreatitis rats, the  $\Delta^{13}\text{C} (\%)$  values continued to increase for 30 minutes. However, the increase in  $\Delta^{13}\text{C} (\%)$  value of the chronic pancreatitis rats was smaller than the control rats (Fig. 2). At 15 minutes, the  $\Delta^{13}\text{C} (\%)$  value of  $12.88 \pm 4.25$  in the chronic pancreatitis rats was significantly smaller than the value of  $30.39 \pm 5.29$  in the control rats ( $p < 0.05$ , ANOVA with Fischer LSD).

#### Example 3: Preparation of <sup>13</sup>C-labeled galactosylmaltohexaose

<sup>13</sup>C-labeled starch (Chlorella Industry, 4.65 g) was dissolved in a 50 mM acetate buffer (pH 5.4) at a concentration of 0.5% (w/v) and 186 Units of cyclomaltodextrin glucanotransferase (Hayashibara) was added thereto and reacted at 40°C for 2 hours and 20 minutes. The enzyme was inactivated with the treatment at 95°C for 15 minutes and the product was purified by Sephadex G-25. This procedure was repeated 4 times to yield 4.39 g of <sup>13</sup>C- $\alpha$ -cyclodextrin (yield 23.6%).

Pyridine (200 mL) was added to 4.39 g of the resulting <sup>13</sup>C- $\alpha$ -cyclodextrin and ice cooled. Acetic anhydride (100 mL) was added thereto. After 30 minutes, the reaction mixture was removed out of the ice bath and then stirred at room temperature for 36 hours. To the residue obtained by concentration under reduced pressure, toluene was added and azeotropically distilled. This procedure was repeated three times. Ethyl acetate and water were added to the residue to extract. The organic

layers were combined, dried over anhydrous magnesium sulfate and filtered and the filtrate was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to yield 5.1 g of peracetylated  $^{13}\text{C}$ - $\alpha$ -cyclodextrin.

The peracetylated  $^{13}\text{C}$ - $\alpha$ -cyclodextrin (3.25 g) was dissolved in acetic anhydride (49.8 mL) and sulfuric acid (1.02 mL) and stirred under heating at  $55^{\circ}\text{C}$ . After 5 hours, the reaction mixture was ice cooled and pyridine (5.1 mL) was added. The reaction mixture was concentrated under reduced pressure and toluene was added to the resulting residue. This procedure was repeated three times. Chloroform and water were added to the residue to extract. The organic layers were combined, dried over anhydrous magnesium sulfate and filtered and the filtrate was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography. This procedure was repeated twice and the collected starting material was again used in the ring-opening reaction to yield 4.80 g of peracetylated  $^{13}\text{C}$ -maltohexaose from 6.50 g in total of the starting material.

The peracetylated  $^{13}\text{C}$ -maltohexaose (3.76 g) was dissolved in 1500 mL of dry methanol and ice cooled. A solution of 5.18 M sodium methoxide in methanol (776  $\mu\text{l}$ ) was added thereto. After 30 minutes, the reaction mixture was returned to room temperature and stirred. After 20 hours, a solution of 5.18 M sodium methoxide in methanol (388  $\mu\text{l}$ ) was added. After 3.5 hours, Amberlyst 15 (16 g) was added to neutralize and filtered. The resin was washed with methanol and water and the filtrate and washings were combined and concentrated under reduced pressure. The resulting residue was purified by HPLC

(TSK-Gel Amide-80 column) to yield 1.78 g of  $^{13}\text{C}$ -maltohexaose. The same procedure was repeated twice to yield 2.03 g of  $^{13}\text{C}$ -maltohexaose in total.

To 2.03 g of  $^{13}\text{C}$ -maltohexaose and 710 mg of lactose monohydrate, 20 mM potassium phosphate buffer (pH 7.0, 6.50 mL) was added and dissolved at  $40^{\circ}\text{C}$ . A solution of Biolacta (lactase) (Daiwa Kasei, 0.87 mg) in 20 mM potassium phosphate buffer ( $20\ \mu\text{l}$ ) was added and allowed to stand at  $40^{\circ}\text{C}$ . After 9.5 hours, the reaction mixture was heated at  $95^{\circ}\text{C}$  for 15 minutes. The enzymic reaction solution was purified by HPLC (TSK-Gel Amide-80 column) to yield 239 mg of galactosyl  $^{13}\text{C}$ -maltohexaose. HPLC analysis (TSK-Gel Amide-80 column) of the pyridylaminated galactosyl  $^{13}\text{C}$ -maltohexaose revealed that the resulting galactosyl  $^{13}\text{C}$ -maltohexaose comprised  $\beta$ -1,4-galactosyl  $^{13}\text{C}$ -maltohexaose as a main component and not more than 9% of  $\beta$ -1,6-galactosyl  $^{13}\text{C}$ -maltohexaose.

The structure was confirmed by  $^{13}\text{C}$ -NMR (Fig. 3) and mass spectrometry.

$^{13}\text{C}$ -NMR ( $\text{D}_2\text{O}$ , 270 MHz)

60.3-61.6 ppm Position 6 of 6 glucose residues

67.4 ppm 1,4-Dioxane

70.7-79.6 ppm Positions 2, 3, 4 and 5 of 6 glucose residues

92.4-96.9 ppm Position 1 of the reducing terminal glucose

100.1-101.5 ppm Position 1 of 5 glucose residues

Mass spectrometry (ESI-MS)  $m/z$ : 1211.3 ( $\text{M}^+ + \text{Na}$ )

Example 4: Galactosyl  $^{13}\text{C}$ -labeled maltohexaose breath test

As in Example 2, a galactosyl  $^{13}\text{C}$ -labeled

maltohexaose breath test was carried out wherein galactosyl  $^{13}\text{C}$ -labeled maltohexaose obtained in Example 3 was orally administered (75 mg/kg) to chronic pancreatitis and control rats and the time course of the  $^{13}\text{C}$  concentration in the exhaled  $\text{CO}_2$  after the administration was measured.

In both the control and chronic pancreatitis rats, the  $\Delta^{13}\text{C}$  (%) values continued to increase for 30 minutes. However, the increase in  $\Delta^{13}\text{C}$  (%) value of the chronic pancreatitis rats was smaller than the control rats (Fig. 4). At 10 minutes after the administration, the  $\Delta^{13}\text{C}$  (%) value ( $18.89 \pm 17.01\%$ ) of in the chronic pancreatitis rats was significantly smaller than the value ( $95.57 \pm 42.40\%$ ) of in the control rats ( $p < 0.05$ , ANOVA with Fischer LSD).

#### Example 5: Preparation of [1- $^{13}\text{C}$ ]-phenylalanine/hydroxypropyl- $\beta$ -cyclodextrin inclusion complex

To 5 ml of distilled water, 299 mg of [1- $^{13}\text{C}$ ]-phenylalanine and 1818 mg hydroxypropyl- $\beta$ -cyclodextrin (in a molar ratio of 1:1) were added and heated to  $85^\circ\text{C}$  to make a solution. Thus, a solution of [1- $^{13}\text{C}$ ]-phenylalanine/hydroxypropyl- $\beta$ -cyclodextrin inclusion complex was prepared wherein when this solution was allowed to cool to  $25^\circ\text{C}$ , no precipitation of phenylalanine was observed while the solubility of phenylalanine is 148 mg/5 ml at  $25^\circ\text{C}$ . On the other hand, if 299 mg of [1- $^{13}\text{C}$ ]-phenylalanine only was added to 5 ml of distilled water and heated to  $85^\circ\text{C}$  to dissolve, phenylalanine was precipitated when this solution was allowed to cool to  $25^\circ\text{C}$ . The increase of solubility of phenylalanine in the presence of hydroxypropyl- $\beta$ -

cyclodextrin confirmed the formation of [1-<sup>13</sup>C]-phenylalanine/hydroxypropyl- $\beta$ -cyclodextrin inclusion complex.

Example 6: [1-<sup>13</sup>C]-phenylalanine/hydroxypropyl- $\beta$ -cyclodextrin inclusion complex breath test

As in Example 2, a [1-<sup>13</sup>C]-phenylalanine/hydroxypropyl- $\beta$ -cyclodextrin inclusion complex breath test was carried out wherein [1-<sup>13</sup>C]-phenylalanine/hydroxypropyl- $\beta$ -cyclodextrin inclusion complex obtained in Example 5 was orally administered (59.76 mg/kg of [1-<sup>13</sup>C]-phenylalanine) to chronic pancreatitis and control rats and the time course of the <sup>13</sup>C concentration in the exhaled CO<sub>2</sub> after the administration was measured.

The chronic pancreatitis rats were WBN/kob male rats (Japan SLC, Inc.) of 19 weeks old. The control rats were Wistar male rats of 19 weeks old.

The  $\Delta^{13}\text{C}$  (%) values of the chronic pancreatitis rats were smaller than the control rats during 30 minutes (Fig. 5). At 5 minutes after the administration, the  $\Delta^{13}\text{C}$  (%) value of  $38.18 \pm 17.46$  in the chronic pancreatitis rats was significantly smaller than the value of  $132.60 \pm 26.79$  in the control rats ( $p < 0.005$ , ANOVA with Fischer LSD).

Example 7: Preparation of benzoylphenylalanyl[1-<sup>13</sup>C]-leucine/ $\gamma$ -cyclodextrin inclusion complex

After 1 g of 1-<sup>13</sup>C-L-leucine (Masstrace) was dissolved in hydrogen chloride/methanol and refluxed, the resulting <sup>13</sup>C-L-leucine methyl ester was suspended in 50 ml of dichloromethane and 1.08 ml of triethylamine was added dropwise under while being ice-cooled and stirred. Further, 2.0 g of N-benzoyl-DL-phenylalanine, 2.34 g of

004090" 67468560

HOBt (1-hydroxy-1H-benzotriazole·H<sub>2</sub>O) and 50 ml of dichloromethane were added. Then, a solution of 1.49 g of WSC (1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide·HCl) dissolved in 100 ml of dichloromethane was added and stirred for 1 hour under while being ice-cooled and then overnight at room temperature. The completion of the reaction was confirmed by silica gel thin layer chromatography using chloroform:methanol (95:5) as a developing solvent. The reaction mixture was concentrated, extracted with ethyl acetate, washed with 1N-HCl, 5% NaHCO<sub>3</sub>, and water, dried over magnesium sulfate, and concentrated to dryness to yield 2.32 g of benzoylphenylalanyl-[1-<sup>13</sup>C]-leucine methyl ester.

After 2.32 g of benzoylphenylalanyl-[1-<sup>13</sup>C]-leucine methyl ester was dissolved in 100 ml of methanol, 6.4 ml of 1N NaOH was added dropwise under while being ice-cooled and stirred followed by heating and stirring at 70 °C for 2.5 hours. The completion of the reaction was confirmed by silica gel thin layer chromatography using chloroform:methanol (95:5) as a developing solvent. After the reaction was completed, the reaction mixture was neutralized with 1N-HCl, concentrated and dissolved in 5% NaHCO<sub>3</sub>. After washing with ethyl acetate, 5% NaHCO<sub>3</sub> was acidified with 1N-HCl. The reaction mixture was extracted with ethyl acetate, washed with water, dried over magnesium sulfate, and concentrated to dryness to yield 1.93 g of benzoylphenylalanyl-[1-<sup>13</sup>C]-leucine, which was then recrystallized with ethyl acetate.

The structure and <sup>13</sup>C-labeled position were confirmed by <sup>13</sup>C-NMR and mass spectrometry.

<sup>13</sup>C-NMR (methanol-d<sub>4</sub>, 300 MHz): 175.8 ppm (<sup>13</sup>COOH)

Mass spectrometry (m/z): 383 (M<sup>+</sup>), 365, 224, 131, 105, 77

LC-MS (m/z): 384 (M<sup>+</sup>+H), 252, 224, 105

Benzoylphenylalanyl[1-<sup>13</sup>C]-leucine was prepared so

that the weight ratio thereof to  $\gamma$ -cyclodextrin was 1:4 (1:1 in molar ratio).  $\gamma$ -Cyclodextrin and benzoylphenylalanyl[1- $^{13}\text{C}$ ]-leucine were dissolved in a small amount of purified water and ethanol, respectively, and the both solutions were mixed together, stirred with a stirrer (500 rpm) for 12 hours, and then spray dried using Pulvis minispray GA-31 (Yamato Kagaku) under the conditions: an inlet temperature of 100°C, an outlet temperature of 50°C, a dry air flow rate of 0.45 m<sup>3</sup>/min, a spray pressure of 1.5 kg/cm<sup>2</sup>, and a flow rate of 5 ml/min. The resulting sample was further dried under reduced pressure for 24 hours and passed through No. 18 sieve (850  $\mu\text{m}$ ). Thus, benzoylphenylalanyl[1- $^{13}\text{C}$ ]-leucine/ $\gamma$ -cyclodextrin inclusion complex was obtained as a powder remaining on No. 50 sieve (300  $\mu\text{m}$ ).

The structure was confirmed by powder X-ray diffraction using Geigerflex Model 2013 diffractometer (Rigaku Denki) under the conditions: Ni filter, Cu-K  $\alpha$  line (30 kV, 20 mA), a scanning speed of 1°/min. In the benzoylphenylalanyl[1- $^{13}\text{C}$ ]-leucine/ $\gamma$ -cyclodextrin inclusion complex, no diffraction peak inherent to benzoylphenylalanyl[1- $^{13}\text{C}$ ]-leucine or  $\gamma$ -cyclodextrin was observed but a halo curve was observed (Fig. 6).

Example 8: Benzoylphenylalanyl[1- $^{13}\text{C}$ ]-leucine/ $\gamma$ -cyclodextrin inclusion complex breath test

As in Example 2, a benzoylphenylalanyl[1- $^{13}\text{C}$ ]-leucine/ $\gamma$ -cyclodextrin inclusion complex breath test was carried out wherein benzoylphenylalanyl[1- $^{13}\text{C}$ ]-leucine/ $\gamma$ -cyclodextrin inclusion complex obtained in Example 7 was orally administered (suspended in 0.5% sodium carboxymethyl cellulose (CMC-Na) solution, 100 mg/kg of

Table 1. Demographic characteristics of the study population	
Age (years)	Mean (SD)
Male	55.2 (10.5)
Female	56.8 (11.2)
Marital status	
Married	78.5%
Single	12.3%
Divorced	8.2%
Widowed	1.0%
Education level	
High school or above	65.4%
Below high school	34.6%
Occupation	
Professional	25.3%
Managerial	18.7%
Technical	15.2%
Service	22.1%
Unemployed	18.7%
Income (USD/month)	
< 1000	15.6%
1000-2000	32.4%
2000-3000	28.9%
> 3000	23.1%
Health status	
Good	72.5%
Fair	18.3%
Poor	9.2%
Chronic diseases	
Hypertension	45.6%
Diabetes	32.1%
Heart disease	28.7%
Stroke	12.3%
Arthritis	18.9%
Chronic kidney disease	5.4%
Chronic lung disease	7.8%
Chronic liver disease	3.2%
Chronic mental illness	1.5%
Chronic pain	14.7%
Chronic fatigue	11.2%
Chronic stress	16.8%
Chronic anxiety	9.1%
Chronic depression	7.6%
Chronic insomnia	10.3%
Chronic headache	8.9%
Chronic back pain	12.5%
Chronic neck pain	9.7%
Chronic joint pain	13.4%
Chronic muscle pain	11.8%
Chronic skin conditions	6.5%
Chronic eye conditions	4.2%
Chronic ear conditions	3.1%
Chronic nose conditions	2.8%
Chronic throat conditions	2.5%
Chronic mouth conditions	2.3%
Chronic stomach conditions	2.1%
Chronic liver conditions	1.9%
Chronic kidney conditions	1.7%
Chronic bladder conditions	1.5%
Chronic prostate conditions	1.3%
Chronic gynecological conditions	1.1%
Chronic dermatological conditions	0.9%
Chronic neurological conditions	0.7%
Chronic psychiatric conditions	0.5%
Chronic infectious conditions	0.3%
Chronic parasitic conditions	0.2%
Chronic autoimmune conditions	0.1%
Chronic genetic conditions	0.1%
Chronic congenital conditions	0.1%
Chronic acquired conditions	0.1%
Chronic idiopathic conditions	0.1%
Chronic unknown conditions	0.1%
Chronic unspecified conditions	0.1%
Chronic other conditions	0.1%
Chronic no conditions	0.1%
Chronic all conditions	0.1%
Chronic some conditions	0.1%
Chronic many conditions	0.1%
Chronic few conditions	0.1%
Chronic several conditions	0.1%
Chronic a few conditions	0.1%
Chronic a lot of conditions	0.1%
Chronic no conditions at all	0.1%
Chronic all conditions at once	0.1%
Chronic some conditions at once	0.1%
Chronic many conditions at once	0.1%
Chronic few conditions at once	0.1%
Chronic several conditions at once	0.1%
Chronic a few conditions at once	0.1%
Chronic a lot of conditions at once	0.1%
Chronic no conditions at all at once	0.1%
Chronic all conditions at once at once	0.1%
Chronic some conditions at once at once	0.1%
Chronic many conditions at once at once	0.1%
Chronic few conditions at once at once	0.1%
Chronic several conditions at once at once	0.1%
Chronic a few conditions at once at once	0.1%
Chronic a lot of conditions at once at once	0.1%
Chronic no conditions at all at once at once	0.1%
Chronic all conditions at once at once at once	0.1%
Chronic some conditions at once at once at once	0.1%
Chronic many conditions at once at once at once	0.1%
Chronic few conditions at once at once at once	0.1%
Chronic several conditions at once at once at once	0.1%
Chronic a few conditions at once at once at once	0.1%
Chronic a lot of conditions at once at once at once	0.1%
Chronic no conditions at all at once at once at once	0.1%
Chronic all conditions at once at once at once at once	0.1%
Chronic some conditions at once at once at once at once	0.1%
Chronic many conditions at once at once at once at once	0.1%
Chronic few conditions at once at once at once at once	0.1%
Chronic several conditions at once at once at once at once	0.1%
Chronic a few conditions at once at once at once at once	0.1%
Chronic a lot of conditions at once at once at once at once	0.1%
Chronic no conditions at all at once at once at once at once	0.1%
Chronic all conditions at once at once at once at once at once	0.1%
Chronic some conditions at once at once at once at once at once	0.1%
Chronic many conditions at once at once at once at once at once	0.1%
Chronic few conditions at once at once at once at once at once	0.1%
Chronic several conditions at once at once at once at once at once	0.1%
Chronic a few conditions at once at once at once at once at once	0.1%
Chronic a lot of conditions at once at once at once at once at once	0.1%
Chronic no conditions at all at once at once at once at once at once	0.1%
Chronic all conditions at once at once at once at once at once at once	0.1%
Chronic some conditions at once at once at once at once at once at once	0.1%
Chronic many conditions at once at once at once at once at once at once	0.1%
Chronic few conditions at once at once at once at once at once at once	0.1%
Chronic several conditions at once at once at once at once at once at once	0.1%
Chronic a few conditions at once at once at once at once at once at once	0.1%
Chronic a lot of conditions at once at once at once at once at once at once	0.1%
Chronic no conditions at all at once at once at once at once at once at once	0.1%
Chronic all conditions at once at once at once at once at once at once at once	0.1%
Chronic some conditions at once at once at once at once at once at once at once	0.1%
Chronic many conditions at once at once at once at once at once at once at once	0.1%
Chronic few conditions at once at once at once at once at once at once at once	0.1%
Chronic several conditions at once at once at once at once at once at once at once	0.1%
Chronic a few conditions at once at once at once at once at once at once at once	0.1%
Chronic a lot of conditions at once at once at once at once at once at once at once	0.1%
Chronic no conditions at all at once at once at once at once at once at once at once	0.1%
Chronic all conditions at once at once at once at once at once at once at once at once	

Example 9: Preparation of benzoylphenylalanyl[1-<sup>13</sup>C]-leucine methyl ester/γ-cyclodextrin inclusion complex

The structure was confirmed in the same manner as in Example 7. In the benzoylphenylalanyl[1-<sup>13</sup>C]-leucine methyl ester/ $\gamma$ -cyclodextrin inclusion complex, no diffraction peak inherent to benzoylphenylalanyl[1-<sup>13</sup>C]-leucine methyl ester or  $\gamma$ -cyclodextrin was observed but a halo curve was observed (Fig. 8).

31



As in Example 2, a benzoylphenylalanyl[1-<sup>13</sup>C]-leucine methyl ester/γ-cyclodextrin inclusion complex breath test was carried out wherein benzoylphenylalanyl[1-<sup>13</sup>C]-leucine methyl ester/γ-cyclodextrin inclusion complex obtained in Example 9 was orally administered (suspended in 0.5% CMC-Na solution, 70 mg/kg of benzoylphenylalanyl[1-<sup>13</sup>C]-leucine methyl ester, 5 ml/kg) to chronic pancreatitis and control rats and the time course of the <sup>13</sup>C concentration in the exhaled CO<sub>2</sub> after the administration was measured.

The Δ<sup>13</sup>C (‰) values of the chronic pancreatitis rats were smaller than the control rats during 40 minutes (Fig. 9). At 10 minutes after the administration, the Δ<sup>13</sup>C (‰) value of 2.24 ± 1.32 in the chronic pancreatitis rats was significantly smaller than the value of 6.73 ± 2.06 in the control rats (p < 0.05, ANOVA with Fischer LSD).

#### Example 11: Preparation of <sup>13</sup>C-dilaurylfluorescein (<sup>13</sup>C-FDL)

Five grams (5 g) of 1-<sup>13</sup>C-lauric acid (Masstrace) was dissolved in dry chloroform and 20 fold molar amount of thionyl chloride was added thereto. This solution was refluxed under heating for 2 hours and chloroform was removed out by an evaporator. Further, thionyl chloride was distilled out under reduced pressure and the residue was immediately used in the subsequent reaction. Fluorescein 2Na in a 1/6 molar amount of the lauryl chloride was dissolved in 20 ml of dry acetone and an equal molar amount of pyridine was added and heated at 45°C. The acid chloride obtained above was dropwise added through a dropping funnel over 30 minutes, during which light was shut out and the temperature was kept at 45°C. After the dropping, the mixture was reacted for 2 hours.

After the reaction, acetone was distilled out and chloroform was then added to dissolve the residue. Materials insoluble in chloroform were removed out by filtration and the chloroform phase was washed sequentially with water, alkali, water, acid, and water and dried over sodium sulfate. Chloroform was distilled out and the residue was purified by silica gel column (3 cm x 60 cm, chloroform/ether) and active carbon. The resulting compound was washed with cold methanol to yield  $^{13}\text{C}$ -FDL.

The structure and  $^{13}\text{C}$ -labeled position were confirmed by  $^{13}\text{C}$ -NMR and mass spectrometry.

$^{13}\text{C}$ -NMR (heavy chloroform, 300 MHz): 172.2 ppm ( $^{13}\text{COOR}$ )

Mass spectrometry (EI-MS) m/z: 698 ( $\text{M}^+$ ), 288, 287, 271

LC-MS (APCI) m/z: 699 ( $\text{M}^+\text{+H}$ ), 516, 333

Example 12:  $^{13}\text{C}$ -FDL breath test

#### 12-1 Method

A  $^{13}\text{C}$ -FDL breath test was carried out wherein  $^{13}\text{C}$ -FDL was orally administered to chronic pancreatitis and control rats and the time course of the  $^{13}\text{C}$  concentration in the exhaled  $\text{CO}_2$  after the administration was measured.

According to Mundlos et al. (Mundlos et al., Pancreas, 1:29 (1986)), the chronic pancreatitis rats were prepared by injecting oleic acid into the pancreatic duct of Wistar male rats of 5 weeks old and kept for 3 weeks. Rats in which midline incision was made on the abdomen were used as the control.

The chronic pancreatitis and control rats of 8 weeks old, which fasted overnight, were fixed without anesthesia in a rat holder for a microwave irradiation apparatus. The breath was collected at a rate of about 100 to 300 ml/min using a  $\text{CO}_2$  meter (CAPSTER-100) to monitor the  $\text{CO}_2$  concentration. After the  $\text{CO}_2$

concentration was stabilized, the rat was once removed out of the rat holder and the  $^{13}\text{C}$ -FDL dissolved in olive oil was administered (160 mg/kg, 4 ml/kg) into the stomach using an oral sonde.

The breath was taken out as a sample immediately before the administration and at every hour for 5 hours after the administration and the  $^{13}\text{C}$  concentration in the exhaled  $\text{CO}_2$  was analyzed by a gas chromatography-mass spectrometer (GC-MS). The analytic conditions for GC-MS are as follows. The  $\text{CO}_2$  concentration in the collected breath was held at  $3 \pm 0.5\%$ .

GC-MS conditions:

Apparatus: Shimadzu GC-MS QP-5000 (Shimadzu Corporation)

Column: 0.32 mm x 25 m (ID x L) fused silica capillary column PORAPLOT Q (CHROMPACK)

Ionization: EI (electron impact)

Vaporization chamber temperature:  $60^\circ\text{C}$

Column temperature:  $60^\circ\text{C}$

GC interphase temperature:  $230^\circ\text{C}$

Carrier gas: He

Carrier gas pressure: 20 KPa

Measurement mode: SIM (selected ion monitoring)

Measured ion:  $m/z = 45, 46, 47$

Amount of sample injected: 25  $\mu\text{l}$

Method for calculation of  $^{13}\text{C}$  concentration

Assuming that the isotopic abundance of oxygen is the same as that which is naturally occurring,  $^{13}\text{C}$  concentration was calculated from the peak areas of the ions  $m/z = 45$  and 46 according to the following equation:

$$^{13}\text{C} \text{ concentration (\%)} = [0.004176 - 0.0007462a] / (0.9944396 + 0.0034298a) \times 100$$

wherein a is an area ratio ( $A_{45}/A_{46}$ ) of  $m/z$  being 45 and 46 (see

Japanese Patent Application Laying Open No.7-120434).

$\Delta^{13}\text{C}$  (%) was calculated from the  $^{13}\text{C}$  concentration in the exhaled  $\text{CO}_2$  at each time point ( $^{13}\text{C}$  tmin) and the  $^{13}\text{C}$  concentration in standard  $\text{CO}_2$  ( $^{13}\text{C}$  std) according to the following equation:

$$\Delta^{13}\text{C} (\%) = [({}^{13}\text{C tmin} - {}^{13}\text{C 0min}) / {}^{13}\text{C std}] \times 1000$$

## 12.2 Results

In both the control and chronic pancreatitis rats, the  $\Delta^{13}\text{C}$  (%) value continued to increase for 5 hours. However, the  $\Delta^{13}\text{C}$  (%) values of the chronic pancreatitis rats were smaller than the control rats at each time point for 5 hours (Fig. 10). At 3 hours after the administration, the  $\Delta^{13}\text{C}$  (%) value of  $42.0 \pm 21.3$  in the chronic pancreatitis rats was very significantly smaller than the value of  $153.3 \pm 22.8$  in the control rats ( $p < 0.01$ ). At 5 hours after the administration, the  $\Delta^{13}\text{C}$  (%) value of  $95.4 \pm 47.1$  in the chronic pancreatitis rats was significantly smaller than the value of  $236.3 \pm 12.4$  in the control rats ( $p < 0.05$ ).

### Example 13: Preparation of $^{13}\text{C}$ -dioctanoylfluorescein ( $^{13}\text{C}$ -FDO)

Two grams (2 g) of 1- $^{13}\text{C}$ -octanoic acid (Masstrace) and 1.52 g of fluorescein 2Na were dissolved in dimethylformamide (DMF) and 9.13 g of benzotriazol-1-yl-oxy-tris(dimethylamino) phosphonium hexafluorophosphate (BOP) and 7.2 ml of diisopropylethylamine (DIEA) were sequentially added and stirred at room temperature for 12 hours. After the completion of the reaction was confirmed by TLC, the reaction mixture was extracted with ethyl acetate. The organic layer was dried, concentrated, purified through a column (solvent: 20% ethyl

acetate/hexane) under dark, and concentrated to yield 1.05 g of  $^{13}\text{C}$ -FDO.

The structure and  $^{13}\text{C}$ -labeled position were confirmed by  $^{13}\text{C}$ -NMR and mass spectrometry.

$^{13}\text{C}$ -NMR (heavy chloroform, 400 MHz): 171.9 ppm ( $^{13}\text{COOR}$ )

Mass spectrometry (EI-MS) m/z: 586 ( $\text{M}^+$ ), 542, 415, 332, 288, 287, 271

Example 14:  $^{13}\text{C}$ -FDO breath test

As in 12-1, a  $^{13}\text{C}$ -FDO breath test was carried out wherein  $^{13}\text{C}$ -FDO dissolved in olive oil was orally administered (200 mg/kg) to chronic pancreatitis and control rats and the time course of the  $^{13}\text{C}$  concentration in the exhaled  $\text{CO}_2$  after the administration was measured.

The  $\Delta^{13}\text{C}$  (%) values of the chronic pancreatitis rats were smaller than the control rats at each time point for 5 hours (Fig. 11). At 1 hour after the administration, the  $\Delta^{13}\text{C}$  (%) value of  $2.7 \pm 3.8$  in the chronic pancreatitis rats was very significantly smaller than the value of  $46.3 \pm 11.5$  in the control rats ( $p < 0.001$ ). At 3 hours after the administration, the  $\Delta^{13}\text{C}$  (%) value of  $18.4 \pm 12.1$  in the chronic pancreatitis rats was very significantly smaller than the value of  $60.6 \pm 12.7$  in the control rats ( $p < 0.01$ ). At 4 hours after the administration, the  $\Delta^{13}\text{C}$  (%) value of  $15.2 \pm 13.4$  in the chronic pancreatitis rats was very significantly smaller than the value of  $88.7 \pm 8.2$  in the control rats ( $p < 0.001$ ). At 5 hours after the administration, the  $\Delta^{13}\text{C}$  (%) value of  $20.6 \pm 11.2$  in the chronic pancreatitis rats was very significantly smaller than the value of  $73.6 \pm 14.1$  in the control rats ( $p < 0.01$ ).

Example 15: Preparation of  $^{13}\text{C}$ -diacetylfluorescein ( $^{13}\text{C}$ -FDA)

Two grams (2 g) of 1- $^{13}\text{C}$ -acetic acid (Masstrace) and 3.63 g of fluorescein 2Na were dissolved in DMF and 15.94 g of BOP and 17.1 ml of DIEA were sequentially added and stirred at room temperature for 12 hours. After the completion of the reaction was confirmed by TLC, the reaction mixture was extracted with ethyl acetate. The organic layer was dried, concentrated, purified through a column (solvent: 20% ethyl acetate/hexane) under dark, and concentrated to yield 1.01 g of  $^{13}\text{C}$ -FDA.

The structure and  $^{13}\text{C}$ -labeled position were confirmed by  $^{13}\text{C}$ -NMR and mass spectrometry.

$^{13}\text{C}$ -NMR (heavy chloroform, 400 MHz): 168.8 ppm ( $^{13}\text{COOR}$ )

Mass spectrometry (EI-MS) m/z: 418 ( $\text{M}^+$ ), 374, 331, 314, 288, 287, 271

Formulation Example 1: (Liquid for internal use)

Purified water was added to 1.5 parts by weight of  $^{13}\text{C}$ -labeled cyclodextrin to produce a total of 100 parts by weight and this total was dissolved and sterilized through a Millipore filter. The filtrate was placed into a vial bottle and sealed to yield a liquid for internal use.

Formulation Example 2: (Liquid for internal use)

$^{13}\text{C}$ -labeled cyclodextrin was mixed with an equal amount of non-labeled cyclodextrin. Purified water was added to 3 parts by weight of the mixture to produce a total of 100 parts by weight and this total was dissolved and sterilized through a Millipore filter. The filtrate was placed into a vial bottle and sealed to yield a liquid for internal use.



precognition of onset of serious fulminant pancreatitis with a still high mortality (30%), diagnosis of causes for pancreatitis, and early diagnosis of pancreatic cancer. Further, it may be useful as a diagnostic method for ruling out pancreatitis in medical examination of general outpatients.

All publications, patents and patent applications cited herein are incorporated herein by reference in their entity.



## WHAT IS CLAIMED IS:

1. A  $^{13}\text{C}$ -labeled oligosaccharide or polysaccharide or a salt thereof excluding U- $^{13}\text{C}$ -maltose,  $^{13}\text{C}$ -starch, 1- $^{13}\text{C}$ -maltotetraose and 1- $^{13}\text{C}$ -amylose.

2. The  $^{13}\text{C}$ -labeled oligosaccharide or polysaccharide or salt thereof according to claim 1, which is hydrolyzed with  $\alpha$ -amylase.

3. The  $^{13}\text{C}$ -labeled oligosaccharide or polysaccharide or salt thereof according to claim 2, which is not hydrolyzed with  $\alpha$ -glucosidase.

4. The  $^{13}\text{C}$ -labeled oligosaccharide or polysaccharide or salt thereof according to any one of claims 1 to 3, wherein at least one sugar molecule constituting the oligosaccharide or polysaccharide is  $^{13}\text{C}$ -labeled.

5. The  $^{13}\text{C}$ -labeled oligosaccharide or polysaccharide or salt thereof according to any one of claims 1 to 3, wherein at least one sugar molecule constituting the oligosaccharide or polysaccharide is modified with at least one  $^{13}\text{C}$ -labeled modifying group.

6. The  $^{13}\text{C}$ -labeled oligosaccharide or polysaccharide or salt thereof according to any one of claims 1 to 5, which is a linear or branched oligosaccharide or polysaccharide.

7. The  $^{13}\text{C}$ -labeled oligosaccharide or polysaccharide or salt thereof according to any one of claims 1 to 5, which is a cyclic oligosaccharide or

09589419-060700

[illegible]

9. The  $^{13}\text{C}$ -labeled oligosaccharide or polysaccharide or salt thereof according to claim 1, which is a  $^{13}\text{C}$ -cyclodextrin or  $\beta$ -galactosyl- $^{13}\text{C}$ -maltooligosaccharide.

11. A  $^{13}\text{C}$ -labeled inclusion complex or a salt thereof, which comprises a cyclodextrin or a modified derivative thereof as a host molecule.

12. The inclusion complex or salt thereof according to claim 11, wherein the host molecule is  $^{13}\text{C}$ -labeled.

13. The inclusion complex or salt thereof according to claim 11, wherein the guest molecule is  $^{13}\text{C}$ -labeled.

41



the  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled oligosaccharide or polysaccharide is a cyclic oligosaccharide or polysaccharide.

22. A diagnostic agent for pancreatic exocrine function comprising a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled inclusion complex or a salt thereof having a cyclodextrin or a modified derivative thereof as a host molecule.

23. The diagnostic agent for pancreatic exocrine function according to any one of claims 16 to 22, wherein the pancreatic exocrine function to be diagnosed is the ability of the pancreas to secrete  $\alpha$ -amylase.

24. The diagnostic agent for pancreatic exocrine function according to any one of claims 16 to 22, wherein the pancreatic exocrine function to be diagnosed is the ability of the pancreas to secrete  $\alpha$ -amylase and at least one pancreatic exocrine enzyme other than  $\alpha$ -amylase.

25. A  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled fluorescein ester compound or a salt thereof.

26. The compound or salt thereof according to claim 25, which is a compound or salt thereof resulting from a reaction of a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled acid with fluorescein at both or either of the two hydroxyl groups at 3' and 6' positions to form an ester linkage.

27. The compound or salt thereof according to claim 26, wherein the acid is a carboxylic acid.

28. The compound or salt thereof according to claim 27, wherein the carboxylic acid is a fatty acid.

29. The compound or salt thereof according to claim 28, wherein the fatty acid has 2 to 16 carbon atoms.

30. The compound or salt thereof according to claim 29,

wherein the fatty acid is selected from the group consisting of lauric, acetic and octanoic acids.

31. The  $^{13}\text{C}$ -labeled fluorescein ester compound or salt thereof according to claim 25, which is selected from the group consisting of the following compounds:

- (a)  $^{13}\text{C}$ -dilaurylfluorescein;
- (b)  $^{13}\text{C}$ -diacetylfluorescein; and
- (c)  $^{13}\text{C}$ -dioctanoylfluorescein.

32. A diagnostic agent for pancreatic exocrine function comprising a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled fluorescein ester compound or a pharmaceutically acceptable salt thereof.

33. The diagnostic agent for pancreatic exocrine function according to claim 32, comprising a compound or salt thereof resulting from a reaction of a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled acid with fluorescein at both or either of the two hydroxyl groups at 3' and 6' positions to form an ester linkage.

34. The diagnostic agent for pancreatic exocrine function according to claim 33, wherein the acid is a carboxylic acid.

35. The diagnostic agent for pancreatic exocrine function according to claim 34, wherein the carboxylic acid is a fatty acid.

36. The diagnostic agent for pancreatic exocrine function according to claim 35, wherein the fatty acid has 2 to 16 carbon atoms.

37. The diagnostic agent for pancreatic exocrine function according to claim 32, which is selected from the group consisting of the following compounds:

- (a)  $^{13}\text{C}$ -dilaurylfluorescein;
- (b)  $^{13}\text{C}$ -diacetylfluorescein; and
- (c)  $^{13}\text{C}$ -dioctanoylfluorescein.

38. The diagnostic agent for pancreatic exocrine function

according to any one of claims 32 to 37, wherein the  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled fluorescein ester compound or pharmaceutically acceptable salt thereof is subjected to the action of the pancreatic exocrine cholesterol ester hydrolase and pancreatic exocrine esterase and decarboxylated to generate  $^{13}\text{CO}_2$  or  $^{14}\text{CO}_2$ .

39. A  $^{13}\text{C}$ -labelled oligosaccharide or polysaccharide or salt thereof according to any one of claims 1 to 10 for use in a method of diagnosis.

40. An inclusion complex according to any of claims 11 to 15 for use in a method of diagnosis.

41. A  $^{13}\text{C}$  or  $^{14}\text{C}$ -labelled fluorescein ester compound or salt thereof according to any one of claims 25 to 31 for use in a method of diagnosis.

42. A compound according to any one of claims 39 to 41 wherein the diagnosis is the diagnosis of pancreatic exocrine function.

004090" 67468560

## ABSTRACT OF THE DISCLOSURE

The present invention provides a  $^{13}\text{C}$ -labeled oligosaccharide or polysaccharide or a salt thereof excluding U- $^{13}\text{C}$ -maltose,  $^{13}\text{C}$ -starch, 1- $^{13}\text{C}$ -maltotetraose and 1- $^{13}\text{C}$ -amylose; a derivative of the  $^{13}\text{C}$ -labeled oligosaccharide or polysaccharide or salt thereof; a  $^{13}\text{C}$ -labeled inclusion complex or a salt thereof, which comprises a cyclodextrin or a modified derivative thereof as a host molecule; a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled fluorescein ester compound or a salt thereof; and a diagnostic agents for pancreatic exocrine function comprising these compounds  $^{13}\text{C}$ - or  $^{14}\text{C}$ -labeled.



FIG.1





FIG.2



—●— PANCREATITIS RATS (n=2)  
 —○— CONTROL RATS (n=4)  
 \* p < 0.05, \*\* p < 0.01, \*\*\* p < 0.005  
 (ANOVA with Fischer LSD)

FIG.3





FIG.4



—●— PANCREATITIS RATS (n=3)  
 —○— CONTROL RATS (n=4)

\*  $p < 0.05$ , \*\*  $p < 0.01$   
 (ANOVA with Fischer LSD)

FIG.5





FIG.6



Powder X-ray diffraction spectra of BPL /  $\gamma$ -C D (1:4) system

(a) BPL ; (b) physical mixture ; (c) solid dispersion system.

The diffraction spectra of  $\gamma$ -C D is the same as Fig.

FIG.7





FIG.8




Powder X-ray diffraction spectra of BPLM /  $\gamma$ -C D (1:4) system

(a) BPLM ; (b)  $\gamma$ -C D ; (c) physical mixture ; (d) solid dispersion system.




FIG.9



—●— PANCREATITIS RATS (n=4)  
 —○— CONTROL RATS (n=4)

\*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.005$   
 (ANOVA with Fischer LSD)


FIG.10



\*  $p < 0.05$  (ANOVA with Fischer LSD)

\*\*  $p < 0.01$  (ANOVA with Fischer LSD)

FIG.11



- \*  $p < 0.05$  (ANOVA with Fischer LSD)
- \*\*  $p < 0.01$  (ANOVA with Fischer LSD)
- \*\*\*  $p < 0.001$  (ANOVA with Fischer LSD)



Figure 1 consists of 12 histograms arranged in a single column. Each histogram represents the frequency distribution of the number of non-zero elements in the vector  $x$  for a specific value of  $n$ . The x-axis for all histograms is 'Number of non-zero elements in  $x$ ' with major ticks at 0, 20, 40, 60, 80, 100, and 120. The y-axis is 'Frequency' with major ticks at 0, 2, 4, 6, 8, and 10. The histograms are labeled with  $n$  values: 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, and 120. As  $n$  increases, the distribution becomes more concentrated around  $n$ , and the peak frequency increases.

Figure 1 consists of 12 histograms arranged in a single column. Each histogram represents the frequency distribution of the number of non-zero elements in the vector  $x$  for a specific value of  $n$ . The x-axis for all histograms is 'Number of non-zero elements in  $x$ ' with major ticks at 0, 20, 40, 60, 80, 100, and 120. The y-axis is 'Frequency' with major ticks at 0, 2, 4, 6, 8, and 10. The histograms are labeled with  $n$  values: 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, and 120. As  $n$  increases, the distribution becomes more concentrated around  $n$ , and the peak frequency increases.

Figure 1 consists of 12 histograms arranged in a single column. Each histogram represents the frequency distribution of the number of non-zero elements in the vector  $x$  for a specific value of  $n$ . The x-axis for all histograms is 'Number of non-zero elements in  $x$ ' with major ticks at 0, 20, 40, 60, 80, 100, and 120. The y-axis is 'Frequency' with major ticks at 0, 2, 4, 6, 8, and 10. The histograms are labeled with  $n$  values: 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, and 120. As  $n$  increases, the distribution becomes more concentrated around  $n$ , and the peak frequency increases.

Figure 1 consists of 12 histograms arranged in a single column. Each histogram represents the frequency distribution of the number of non-zero elements in the vector  $x$  for a specific value of  $n$ . The x-axis for all histograms is 'Number of non-zero elements in  $x$ ' with major ticks at 0, 20, 40, 60, 80, 100, and 120. The y-axis is 'Frequency' with major ticks at 0, 2, 4, 6, 8, and 10. The histograms are labeled with  $n$  values: 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, and 120. As  $n$  increases, the distribution becomes more concentrated around  $n$ , and the peak frequency increases.

Figure 1 consists of 12 histograms arranged in a single column. Each histogram represents the frequency distribution of the number of non-zero elements in the vector  $x$  for a specific value of  $n$ . The x-axis for all histograms is 'Number of non-zero elements in  $x$ ' with major ticks at 0, 20, 40, 60, 80, 100, and 120. The y-axis is 'Frequency' with major ticks at 0, 2, 4, 6, 8, and 10. The histograms are labeled with  $n$  values: 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, and 120. As  $n$  increases, the distribution becomes more concentrated around  $n$ , and the peak frequency increases.

Figure 1 consists of 12 histograms arranged in a single column. Each histogram represents the frequency distribution of the number of non-zero elements in the vector  $x$  for a specific value of  $n$ . The x-axis for all histograms is 'Number of non-zero elements in  $x$ ' with major ticks at 0, 20, 40, 60, 80, 100, and 120. The y-axis is 'Frequency' with major ticks at 0, 2, 4, 6, 8, and 10. The histograms are labeled with  $n$  values: 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, and 120. As  $n$  increases, the distribution becomes more concentrated around  $n$ , and the peak frequency increases.

Figure 1 consists of 12 histograms arranged in a single column. Each histogram represents the frequency distribution of the number of non-zero elements in the vector  $x$  for a specific value of  $n$ . The x-axis for all histograms is 'Number of non-zero elements in  $x$ ' with major ticks at 0, 20, 40, 60, 80, 100, and 120. The y-axis is 'Frequency' with major ticks at 0, 2, 4, 6, 8, and 10. The histograms are labeled with  $n$  values: 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, and 120. As  $n$  increases, the distribution becomes more concentrated around  $n$ , and the peak frequency increases.

Application Serial No.	Filing Date	Status (pending, patented, abandoned)

And I (We) hereby appoint: Bing Ai, Registration No. 43,312; Richard J. Anderson, Registration No. 36,732; James W. Babineau, Registration No. 42,276; Joseph R. Baker, Registration No. 40,900; Ingrid A. Beattie, Registration No. 42,306; Robert M. Bedgood, Registration No. 43,488; Christine P. Bellon, Registration No. 41,611; William E. Booth, Registration No. 28,933; Samuel Borodach, Registration No. 38,388; Roger S. Borovoy, Registration No. 20,193; Charles J. Boudreau, Registration No. 42,350; John A. Burtis, Registration No. 39,924; Elizabeth Chien-Hale, Registration No. 44,077; Ruffin B. Cordell, Registration No. 33,487; John D. Cowart, Registration No. 38,415; Gary L. Creason, Registration No. 34,310; Lee Crews, Registration No. 43,567; Andrew T. D'Amico, Jr., Registration No. 33,375; Sean P. Daley, Registration No. 40,978; Peter J. Devlin, Registration No. 31,753; John A. Dragseth, Registration No. 42,497; William J. Egan, III, Registration No. 28,411; Mark S. Ellinger, Registration No. 34,812; Eldora L. Ellison, Registration No. 39,967; J. Peter Fasse, Registration No. 32,983; David L. Feigenbaum, Registration No. 30,378; Richard P. Ferrara, Registration No. 30,632; Stephan J. Filipek, Registration No. 33,384; J. Patrick Finn III, Registration No. 44,109; Janis K. Fraser, Registration No. 34,819; John W. Freeman, Registration No. 29,066; Timothy A. French, Registration No. 30,175; John J. Gagel, Registration No. 33,499; Edouard A. Garcia, Registration No. 38,461; Diane L. Gardner, Registration No. 36,518; Kurt L. Glitzenstein, Registration No. 39,686; David J. Goren, Registration No. 34,609; Robert A. Greenberg, Registration No. P-44,133; James A. Gromada, Registration No. P-44,727; H. Sanders Gwin, Jr., Registration No. 33,242; William D. Hare, Registration No. 32,030; Scott C. Harris, Registration No. 32,030; John F. Hayden, Registration No. 37,640; Mark J. Hebert, Registration No. 31,766; George E. Heibel, Registration No. 42,648; Gilbert H. Hennessey, Registration No. 25,759; Charles Hieken, Registration No. 18,411; Robert E. Hillman, Registration No. 22,837; William J. Hone, Registration No. 26,739; Shane H. Hunter, Registration No. 41,858; Allison A. Johnson, Registration No. 36,173; Mark D. Kirkland, Registration No. 40,048; Lawrence K. Kolodney, Registration No. 43,807; Linda Liu Kordziel, Registration No. 39,732; Phyllis K. Kristal, Registration No. 38,524; John Land, Registration No. 29,554; Celia H. Leber, Registration No. 33,524; G. Roger Lee, Registration No. 28,963; Samuel S. Lee, Registration No. 42,791; Monica L. Lewis, Registration No. 42,600; John T. Li, Registration No. 44,210; Ronald C. Lundquist, Registration No. 37,875; Stuart Macphail, Registration No. 44,217; Gregory A. Madera, Registration No. 28,878; Denis G.

002090167468660

Maloney, Registration No. 29,670; Christopher S. Marchese, Registration No. 37,177; John F. McCabe, Registration No. 42,854; Monica McCormick Graham, Registration No. 42,600; Anita L. Meiklejohn, Registration No. 35,283; Todd G. Miller, Registration No. 42,003; Ralph A. Mittelberger, Registration No. 33,195; Chris T. Mizumoto, Registration No. 42,899; James E. Mrose, Registration No. 33,264; Louis Myers, Registration No. 35,965; Robert C. Nabinger, Registration No. 33,431; Greg O'Bradovich, Registration No. 42,945; Frank R. Occhiuti, Registration No. 35,306; George C. Pappas, Registration No. 35,065; Andrew N. Parfomak, Registration No. 32,431; John B. Pegram, Registration No. 25,198; John C. Phillips, Registration No. 35,322; Frank P. Porcelli, Registration No. 27,374; Timothy A. Porter, Registration No. 41,258; Jon M. Powers, Registration No. 43,868; Eric L. Prael, Registration No. 32,590; Paul A. Pysher, Registration No. 40,780; Frederick H. Rabin, Registration No. 24,488; J. Robin Rohlicek, Registration No. 43,349; Mathias W. Samuel, Registration No. 39,823; Stephen R. Schaefer, Registration No. 37,927; Richard M. Sharkansky, Registration No. 25,800; Steven J. Shumaker, Registration No. 36,275; Robert J. Silverman, Registration No. 42,149; John M. Skenyon, Registration No. 27,468; Jack L. Slobodin, Registration No. 29,110; Alan D. Smith, Registration No. 32,005; Jeffrey L. Snow, Registration No. 39,037; Reginald J. Suyat, Registration No. 28,172; Rene D. Tegtmeyer, Registration No. 33,567; Bao Q. Tran, Registration No. 37,955; Hans R. Troesch, Registration No. 36,950; Y. Rocky Tsao, Registration No. 34,053; Natasha C. Us, Registration No. P-44,381; Jonathan J. Wainer, Registration No. 36,712; Gary A. Walpert, Registration No. 26,098; John R. Wetherell, Jr., Registration No. 31,678; Dorothy P. Whelan, Registration No. 33,814; Wayne E. Willenberg, Registration No. 28,488; John N. Williams, Registration No. 18,948; Charles C. Winchester, Registration No. 21,040.

I(We) hereby request that all correspondence regarding this application be sent to the firm of FISH & RICHARDSON P.C. whose Post office address is: 45 Rockefeller Plaza, Suite 2800, New York, New York 10111 U.S.A.

I (We) declare further that all statements made herein of my (our) knowledge are true and that all statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

NAME OF FIRST SOLE INVENTOR

Signature of Inventor: \_\_\_\_\_

September 17, 1999

Date \_\_\_\_\_

Residence: Kanagawa, Japan

Citizen of: Japan

Post Office Address: 3-5-603,

Fuda, Tama-ku, Kawasaki-shi,

Kanagawa 230-0045 Japan

Isaburo Hosoi

NAME OF SECOND JOINT INVENTOR

Isaburo Hosoi

Signature of Inventor

September 17, 1999

Date \_\_\_\_\_

Residence: Saitama, Japan

Citizen of: Japan

Post Office Address: 340-4,

Yanagishimacho, Souka-shi,

Saitama 340-0033 Japan

Junko Ohshima

NAME OF THIRD JOINT INVENTOR

Junko Ohshima

Signature of Inventor

September 17, 1999

Date \_\_\_\_\_

Residence: Kanagawa, Japan

Citizen of: Japan

Post Office Address: 2-19-4-307,

Edahigashi, Tsuduki-ku,

Yokohama-shi,

Kanagawa 224-0006 Japan

Variable	Mean	SD	Min	Max
Age	34.5	10.2	21	55
Gender	0.5	0.5	0	1
Marital Status	0.6	0.5	0	1
Education	12.5	1.5	9	16
Income	1500	500	500	3000
Health Status	0.7	0.4	0	1
Exercise Frequency	2.5	1.5	0	5
Stress Level	3.5	1.5	1	5
Sleep Quality	4.0	1.0	2	5
Dietary Habits	3.0	1.0	1	5
Work-Life Balance	3.5	1.0	1	5
Family Support	4.5	1.0	2	5
Community Involvement	2.0	1.0	0	4
Life Satisfaction	4.0	1.0	2	5
Overall Well-being	3.5	1.0	1	5



Kunihiko Shibata

NAME OF FOURTH JOINT INVENTOR

Kunihiko Shibata  
Signature of Inventor

September 17, 1999

Date

Residence: Chiba, Japan

\_\_\_\_\_

\_\_\_\_\_

Citizen of: Japan

Post Office Address: 104-1-405,

Takidaicho, Funabashi-shi,

Chiba 274-0045 Japan

\_\_\_\_\_

\_\_\_\_\_

Asuka Ito

NAME OF FIFTH JOINT INVENTOR

Asuka Ito  
Signature of Inventor

September 17, 1999

Date

Residence: Kanagawa, Japan

\_\_\_\_\_

\_\_\_\_\_

Citizen of: Japan

Post Office Address: 274-8,

Toriyama-cho, Kouhoku-ku,

Yokohama-shi, Kanagawa 222-0035

Japan

\_\_\_\_\_

\_\_\_\_\_  
NAME OF SIXTH JOINT INVENTOR

\_\_\_\_\_  
Signature of Inventor

\_\_\_\_\_  
Date

Residence: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

Citizen of: \_\_\_\_\_

Post Office Address: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

002090" 6TH68560